

# September 12 Math 3260 sec. 53 Fall 2025

## Chapter 2 Systems of Linear Equations

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & & \vdots & & \ddots & & \vdots & = & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array}$$

### 2.2.1 Gaussian Elimination

We're interested in solving such a system using **Gaussian** elimination. Recall that this involved three operations that result in an equivalent system:

- ▶ Multiply an equation by a nonzero scalar  $k$ .  $kE_i \rightarrow E_i$  (scale)
- ▶ Interchange the position of any two equations.  $E_i \leftrightarrow E_j$  (swap)
- ▶ Replace an equation with the sum of itself and a multiple of any other equation.  $kE_i + E_j \rightarrow E_j$  (replace)

## Example

$$\begin{array}{rrcrcl} x_1 & - & x_2 & - & 5x_3 & = & 6 \\ 3x_1 & + & x_2 & - & 7x_3 & = & 10 \\ 2x_1 & - & x_2 & - & 8x_3 & = & 10 \end{array}$$

$$-3E_1 + E_2 \rightarrow E_2$$

$$-2E_1 + E_3 \rightarrow E_3$$

$$x_1 - x_2 - 5x_3 = 6$$

$$4x_2 + 8x_3 = -8$$

$$x_2 + 2x_3 = -2$$

$$\frac{1}{4} E_2 \rightarrow E_2$$

$$-3x_1 + 3x_2 + 15x_3 = -18$$

$$3x_1 + x_2 - 7x_3 = 10$$

$$-2x_1 + 2x_2 + 10x_3 = -12$$

$$2x_1 - x_2 - 8x_3 = 10$$

$$x_1 - x_2 - 5x_3 = 6$$

$$x_2 + 2x_3 = -2$$

$$x_2 + 2x_3 = -2$$

$$-E_2 + E_3 \rightarrow E_3$$

$$x_1 - x_2 - 5x_3 = 6$$

$$x_2 + 2x_3 = -2$$

$$0 = 0$$

we're ready for back substitution.

$$x_2 = \dots - 2x_3$$

$x_3$  is any real number.

$$\begin{aligned}
 x_1 &= 6 + x_2 + 5x_3 \\
 &= 6 + (-2 - 2x_3) + 5x_3 \\
 &= 4 + 3x_3
 \end{aligned}$$

A parametric description of the solution set is, letting  $x_3 = t$

$$x_1 = 4 + 3t$$

$$x_2 = -2 - 2t$$

$$x_3 = t$$

,  $t \in \mathbb{R}$

Converting to vector parametric

$$\vec{x} = \langle x_1, x_2, x_3 \rangle$$

$$= \langle 4+3t, -2-2t, t \rangle$$

$$= \langle 4, -2, 0 \rangle + \langle 3t, -2t, t \rangle$$

$$\vec{x} = \langle 4, -2, 0 \rangle + t \langle 3, -2, 1 \rangle$$

This is a line in  $\mathbb{R}^3$  passing through the point  $(4, -2, 0)$  and parallel to the vector  $\langle 3, -2, 1 \rangle$ .

## Other Solution Cases

An example of the no solution case: The system

$$\begin{array}{rrcrcl} x_1 & + & 4x_2 & + & 3x_3 & = & 1 \\ 2x_1 & + & x_2 & - & x_3 & = & 2 \\ -x_1 & + & 3x_2 & + & 4x_3 & = & 0 \end{array}$$

leads to

$$\begin{array}{rrcrcl} x_1 & + & 4x_2 & + & 3x_3 & = & 1 \\ & & x_2 & + & x_3 & = & 0 \\ & & & & 0 & = & 1 \end{array}$$

**Inconsistent systems** always give rise to an equation that is false.

$$0 = \text{something nonzero}$$

## 2.3 Matrices

### Matrix

A **matrix** (plural *matrices*) is a rectangular array of numbers of the form

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Each number,  $a_{ij}$ , is called an **entry** or an **element** of the matrix. If the matrix has  $m$  rows and  $n$  columns, we say that the **size** or **dimension** of the matrix is “ $m$  by  $n$ ” and write  $m \times n$ .



# Coefficient & Augmented Matrices

Given a system of linear equations with  $m$  equations and  $n$  variables,

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & + & \vdots & + & \ddots & + & \vdots & = & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array},$$

the **coefficient matrix** for the system is the  $m \times n$  matrix

$$\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$



# Coefficient & Augmented Matrices

Given a system of linear equations with  $m$  equations and  $n$  variables,

$$\begin{array}{ccccccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & b_1 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & b_2 \\ \vdots & + & \vdots & + & \ddots & + & \vdots & = & \vdots \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & b_m \end{array},$$

the **augmented matrix** for the system is the  $m \times n + 1$  matrix

$$\left[ \begin{array}{cccc|c} a_{11} & a_{12} & \cdots & a_{1n} & b_1 \\ a_{21} & a_{22} & \cdots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} & b_m \end{array} \right].$$

## Example

Write the coefficient and augmented matrices for each system.

$$\begin{array}{rrcrcl} 2x_1 & + & x_2 & - & 3x_3 & + & x_4 & = & -3 \\ -x_1 & + & 3x_2 & + & 4x_3 & - & 2x_4 & = & 8 \end{array}$$

The coef matrix is  $2 \times 4$

$$\begin{bmatrix} 2 & 1 & -3 & 1 \\ -1 & 3 & 4 & -2 \end{bmatrix}$$

The augmented matrix is  $2 \times 5$

$$\left[ \begin{array}{cccc|c} 2 & 1 & -3 & 1 & -3 \\ -1 & 3 & 4 & -2 & 8 \end{array} \right]$$

## Example

Write the coefficient and augmented matrices for each system.

$$x_1 - 2x_2 + x_3 = 0$$

$$+ 3x_2 - 2x_3 = 0$$

$$x_1 + x_2 - x_3 = 0$$

The coef. matrix is  $3 \times 3$

$$\begin{bmatrix} 1 & -2 & 1 \\ 0 & 3 & -2 \\ 1 & 1 & -1 \end{bmatrix}$$

The augmented matrix is  $3 \times 4$

$$\left[ \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & 3 & -2 & 0 \\ 1 & 1 & -1 & 0 \end{array} \right]$$

# Elementary Row Operations

We will use matrices to perform elimination without involving the symbols and variable names. We have three operations we can perform on a matrix. We'll use the notation

$$R_i$$

to refer to the  $i^{\text{th}}$  row of the matrix.

## Elementary Row Operations

- ▶ Multiply row  $i$  by any nonzero constant  $k$  (**scale**),  $kR_i \rightarrow R_i$ .
- ▶ Interchange row  $i$  and row  $j$  (**swap**),  $R_i \leftrightarrow R_j$ .
- ▶ Replace row  $j$  with the sum of itself and  $k$  times row  $i$  (**replace**),  $kR_i + R_j \rightarrow R_j$ .