

Section 6: Linear Equations Theory and Terminology

We were talking about the basics of linear homogeneous ODEs.

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

And we're assuming that the functions a_1, \dots, a_n are continuous and that $a_n(x) \neq 0$ (at least on some interval I).

The Principle of Superposition

Says that if we have some solutions, say $y_1(x)$, $y_2(x)$, and $y_3(x)$ of a linear homogeneous equation, then every function of the form

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + c_3 y_3(x)$$

is also a solution of that linear, homogeneous equation.

The expression

$$c_1 y_1(x) + c_2 y_2(x) + c_3 y_3(x)$$

is called a **linear combination** of the functions $y_1(x)$, $y_2(x)$, and $y_3(x)$.

We needed a criteria to distinguish functions or characterize their relationship to one another.

Linear Dependence or Independence

Suppose we have a set of functions $f_1(x), f_2(x), \dots, f_n(x)$ defined on some interval I . We can consider the equation

$$c_1 f_1(x) + c_2 f_2(x) + \dots + c_n f_n(x) = 0 \quad \text{for all } x \text{ in } I. \quad (1)$$

Note that it's always possible to pick c 's to make this true (e.g. you can always set all the c values to zero). We'll say that the functions are

- ▶ **Linearly Dependent** if the equation can be made true **with at least one c being nonzero**.
- ▶ **Linearly Independent** if the only way the equation can be true is if all the c 's must be zero.

Determine if the set is Linearly Dependent or Independent on $(-\infty, \infty)$

$$f_1(x) = x^2, \quad f_2(x) = 4x, \quad f_3(x) = x - x^2$$

Can we find c_1, c_2, c_3 such that

$c_1 f_1(x) + c_2 f_2(x) + c_3 f_3(x) = 0$ for all x , with at least one c -nonzero?

Notice that

$$f_3(x) = \frac{1}{4} f_2(x) - f_1(x)$$

$$x - x^2 = \frac{1}{4}(4x) - x^2$$

This can be rearranged (move everything to the left)

as

$$f_1(x) - \frac{1}{4}f_2(x) + f_3(x) = 0$$

This looks like $c_1 f_1 + c_2 f_2 + c_3 f_3 = 0$ with

$$c_1 = 1, \quad c_2 = -\frac{1}{4} \quad \text{and} \quad c_3 = 1.$$

These c 's are not all zero. These

functions are linearly dependent.

Linear Dependence Relation

An equation with at least one c nonzero, such as

$$f_1(x) - \frac{1}{4}f_2(x) + f_3(x) = 0$$

from this last example is called a **linear dependence relation** for the functions $\{f_1, f_2, f_3\}$.

Definition of Wronskian

Let f_1, f_2, \dots, f_n posses at least $n - 1$ continuous derivatives on an interval I . The **Wronskian** of this set of functions is the determinant

$$W(f_1, f_2, \dots, f_n)(x) = \begin{vmatrix} f_1 & f_2 & \cdots & f_n \\ f_1' & f_2' & \cdots & f_n' \\ \vdots & \vdots & \ddots & \vdots \\ f_1^{(n-1)} & f_2^{(n-1)} & \cdots & f_n^{(n-1)} \end{vmatrix}.$$

(Note that, in general, this Wronskian is a function of the independent variable x .)

Determinants

If A is a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then its determinant

$$\det(A) = ad - bc.$$

If A is a 3×3 matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$, then its determinant

$$\det(A) = a_{11}\det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12}\det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13}\det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

Determine the Wronskian of the Functions

$$f_1(x) = \sin x, \quad f_2(x) = \cos x$$

$$W(f_1, f_2)(x) = \begin{vmatrix} f_1 & f_2 \\ f_1' & f_2' \end{vmatrix} \quad \begin{array}{l} f_1'(x) = \cos x \\ f_2'(x) = -\sin x \end{array}$$

$$= \begin{vmatrix} \sin x & \cos x \\ \cos x & -\sin x \end{vmatrix}$$

$$= \sin x (-\sin x) - \cos x (\cos x)$$

$$= -\sin^2 x - \cos^2 x$$

$$= -1$$

$$W(\sin v, \cos x)(x) = -1$$

Determine the Wronskian of the Functions

$$f_1(x) = x^2, \quad f_2(x) = 4x, \quad f_3(x) = x - x^2$$

$$W(f_1, f_2, f_3)(x) = \begin{vmatrix} f_1 & f_2 & f_3 \\ f_1' & f_2' & f_3' \\ f_1'' & f_2'' & f_3'' \end{vmatrix}$$

$$= \begin{vmatrix} x^2 & 4x & x - x^2 \\ 2x & 4 & 1 - 2x \\ 2 & 0 & -2 \end{vmatrix}$$

$$= x^2 \begin{vmatrix} 4 & 1-2x \\ 0 & -2 \end{vmatrix} - 4x \begin{vmatrix} 2x & 1-2x \\ 2 & -2 \end{vmatrix} + (x-x^2) \begin{vmatrix} 2x & 4 \\ 2 & 0 \end{vmatrix}$$

$$= x^2 (-8) - 4x \left(-4x - \underset{-2}{2(1-2x)} \right) + (x-x^2) (-8)$$

$-4x - 2 + 4x$

$$= -8x^2 + 8x - 8x + 8x^2$$

$$= 0$$

$$W(x^2, 4x, x-x^2)(x) = 0$$

Theorem (a test for linear independence)

Let f_1, f_2, \dots, f_n be $n - 1$ times continuously differentiable on an interval I . If there exists x_0 in I such that $W(f_1, f_2, \dots, f_n)(x_0) \neq 0$, then the functions are **linearly independent** on I .

If y_1, y_2, \dots, y_n are n solutions of the linear homogeneous n^{th} order equation on an interval I , then the solutions are **linearly independent** on I if and only if $W(y_1, y_2, \dots, y_n)(x) \neq 0$ for¹ each x in I .

Compute W

If $W = 0 \Rightarrow$ dependent

If $W \neq 0 \Rightarrow$ independent.

¹For solutions of one linear homogeneous ODE, the Wronskian is either always zero or is never zero.

Determine if the functions are linearly dependent or independent:

$$y_1 = e^x, \quad y_2 = e^{-2x} \quad I = (-\infty, \infty)$$

We can use the Wronskian.

$$\begin{aligned} W(y_1, y_2)(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \\ &= \begin{vmatrix} e^x & e^{-2x} \\ e^x & -2e^{-2x} \end{vmatrix} \\ &= e^x (-2e^{-2x}) - e^x (e^{-2x}) \end{aligned}$$

$$= -2e^{-x} - e^{-x} = -3e^{-x}$$

$$W(y_1, y_2)(x) = -3e^{-x}$$

Since W is not just zero,

y_1 and y_2 are linearly
independent.

Fundamental Solution Set

We're still considering this equation

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = 0$$

with the assumptions $a_n(x) \neq 0$ and $a_i(x)$ are continuous on I .

Definition: A set of functions y_1, y_2, \dots, y_n is a **fundamental solution set** of the n^{th} order homogeneous equation provided they

- (i) are solutions of the equation,
- (ii) there are n of them, and
- (iii) they are linearly independent.

Theorem: Under the assumed conditions, the equation has a fundamental solution set.

General Solution of n^{th} order Linear Homogeneous Equation

Let y_1, y_2, \dots, y_n be a fundamental solution set of the n^{th} order linear homogeneous equation. Then the **general solution** of the equation is

$$y(x) = c_1 y_1(x) + c_2 y_2(x) + \cdots + c_n y_n(x),$$

where c_1, c_2, \dots, c_n are arbitrary constants.

Example

Verify that $y_1 = x^2$ and $y_2 = x^3$ form a fundamental solution set of the ODE

$$x^2 y'' - 4xy' + 6y = 0 \quad \text{on } (0, \infty),$$

and determine the general solution.

The ODE is 2nd order. We need to show our two functions are linearly independent solutions.

Are y_1 + y_2 solutions?

$$y_1 = x^2$$

$$y_1' = 2x$$

$$y_1'' = 2$$

$$y_2 = x^3$$

$$y_2' = 3x^2$$

$$y_2'' = 6x$$

Substitute

$$x^2 y_1'' - 4x y_1' + 6y_1 \stackrel{?}{=} 0$$

$$x^2(2) - 4x(2x) + 6x^2 \stackrel{?}{=} 0$$

$$2x^2 - 8x^2 + 6x^2 \stackrel{?}{=} 0$$
$$0 = 0$$

$$x^2 y_2'' - 4x y_2' + 6y_2 \stackrel{?}{=} 0$$

$$x^2(6x) - 4x(3x^2) + 6x^3 \stackrel{?}{=} 0$$

$$6x^3 - 12x^3 + 6x^3 \stackrel{?}{=} 0$$
$$0 = 0$$

y_1 and y_2 are solutions.

We can compute the Wronskian

$$W(y_1, y_2)(x) = \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix}$$

$$= x^2(3x^2) - 2x(x^3)$$

$$= 3x^4 - 2x^4 = x^4$$

$W \neq 0$, they are linearly independent.

The general solution $y = C_1 y_1 + C_2 y_2$

$$y = C_1 x^2 + C_2 x^3.$$