

## Section 7: Reduction of Order

- ▶ We start with a second order, linear, homogeneous ODE in standard form

$$\frac{d^2y}{dx^2} + P(x)\frac{dy}{dx} + Q(x)y = 0.$$

- ▶ We know one solution  $y_1$ . (Keep in mind that  $y_1$  is a known!)
- ▶ We know there is a second linearly independent solution (section 6 theory says so).
- ▶ We try to find  $y_2$  by guessing that it can be found in the form

$$y_2(x) = u(x)y_1(x)$$

where the goal becomes finding  $u$ .

- ▶ **Due to linear independence, we know that  $u$  cannot be constant.**

## Generalization

Consider the equation **in standard form** with one known solution.  
Determine a second linearly independent solution.

$$\frac{d^2 y}{dx^2} + P(x) \frac{dy}{dx} + Q(x)y = 0, \quad y_1(x) \text{ — is known.}$$

Suppose  $y_2 = u y_1$  ← this guess is an ansatz

$$y_2' = u' y_1 + u y_1'$$

$$y_2'' = u'' y_1 + u' y_1' + u' y_1' + u y_1''$$

$$y_2'' = u'' y_1 + 2u' y_1' + u y_1''$$

We know that  $y_1'' + P(x)y_1' + Q(x)y_1 = 0$ .

$$y_2'' + P(x)y_2' + Q(x)y_2 = 0$$

$$\underline{u''}y_1 + z\underline{u'y_1'} + \underline{u y_1''} + P(x)(\underline{u'y_1} + \underline{u y_1'}) + \underline{Q(x)u y_1} = 0$$

Collect  $u''$ ,  $u'$ , and  $u$

$$y_1 u'' + (z y_1' + P(x) y_1) u' + \underbrace{(y_1'' + P(x) y_1' + Q(x) y_1)}_{=0} u = 0$$

since  $y_1$  solves the  
ODE

$u$  solves

$$y_1 u'' + (z y_1' + P(x) y_1) u' = 0$$

Let  $w = u'$ , then  $w' = u''$  and  $w$  solve

$$y_1 w' + (2y_1' + P(x)y_1)w = 0$$

Assume  $y_1(x) \neq 0$  on the domain, and assume  $w > 0$ .

$$w' + \left(2 \frac{y_1'}{y_1} + P(x)\right)w = 0$$

1st order linear and separable. Let's separate variables

$$\frac{dw}{dx} = - \left(2 \frac{dy_1}{dx} \frac{1}{y_1} + P(x)\right)w$$

$$\frac{1}{w} \frac{dw}{dx} = - \left( 2 \frac{\frac{dy_1}{dx}}{y_1} + P(x) \right)$$

$$\frac{1}{w} \frac{dw}{dx} dx = - \left( 2 \frac{dy_1/dx}{y_1} + P(x) \right) dx$$

$$\frac{1}{w} dw = - 2 \frac{dy_1}{y_1} dx - P(x) dx$$

$$\frac{1}{w} dw = - 2 \frac{dy_1}{y_1} - P(x) dx$$

$$\int \frac{1}{w} dw = - 2 \int \frac{dy_1}{y_1} - \int P(x) dx$$

$$\ln w = - 2 \ln |y_1| - \int P(x) dx$$

$$\ln w = \ln y_1^{-2} - \int P(x) dx$$

Exponentiate

$$\begin{aligned} w &= e^{\ln y_1^{-2} - \int P(x) dx} \\ &= e^{\ln y_1^{-2}} \cdot e^{-\int P(x) dx} \end{aligned}$$

$$\Rightarrow w = \frac{e^{-\int P(x) dx}}{y_1^2}$$

$$w = u' \Rightarrow u = \int w dx$$

Hence

$$u = \int \frac{e^{-\int p(x) dx}}{y_1^2} dx$$

$y_2 = uy_1$  and the general solution

$$y = C_1 y_1 + C_2 y_2$$

## Reduction of Order Formula

For the second order, homogeneous equation **in standard form** with one known solution  $y_1$ , a second linearly independent solution  $y_2$  is given by

$$y_2 = y_1(x) \int \frac{e^{-\int P(x) dx}}{(y_1(x))^2} dx$$



## Example

Find the solution of the IVP where one solution of the ODE is given.

$$y'' + 4y' + 4y = 0 \quad y_1 = e^{-2x}, \quad y(0) = 1, \quad y'(0) = 1$$

Find the general solution  $y = C_1 y_1 + C_2 y_2$

We can use reduction of order to find  $y_2$ .

$$y_2 = u y_1 \quad \text{where} \quad u = \int \frac{e^{-\int p(x) dx}}{y_1^2} dx$$

The ODE is in standard form  $P(x) = 4$

$$-\int p(x)dx = -\int 4dx = -4x \quad (\text{ignoring } +C)$$

$$e^{\int p(x)dx} = e^{-4x}. \quad \text{Given } y_1 = e^{-2x}$$

$$u = \int \frac{e^{\int p(x)dx}}{y_1^2} dx = \int \frac{e^{-4x}}{(e^{-2x})^2} dx$$

$$= \int \frac{e^{-4x}}{e^{-4x}} dx = \int dx = x$$

$$y_2 = uy_1 = x e^{-2x}$$

The general solution to the ODE is

$$y = c_1 e^{-2x} + c_2 x e^{-2x}$$

Apply  $y(0) = 1$  ,  $y'(0) = 1$

$$y' = -2c_1 e^{-2x} + c_2 e^{-2x} - 2c_2 x e^{-2x}$$

$$y(0) = c_1 e^0 + c_2 \cdot 0 \cdot e^0 = 1 \Rightarrow c_1 = 1$$

$$y'(0) = -2c_1 e^0 + c_2 e^0 - 2c_2 \cdot 0 \cdot e^0 = 1$$

$$-2c_1 + c_2 = 1 \Rightarrow c_2 = 1 + 2c_1 = 1 + 2 = 3$$

The solution to the IVP is  $y = e^{-2x} + 3x e^{-2x}$