

Question: Could the method of undetermined coefficients be used to find particular solutions to the ODEs

$$y'' + y = \tan x, \quad \text{or} \quad x^2 y'' + xy' - 4y = e^x?$$

(Why or why not?)



this side
right is
not the
right type



this left is
not constant
coefficient.

Section 10: Variation of Parameters

We're still interested in linear, nonhomogeneous ODEs. Something of the form

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x).$$

The previous method (undetermined coefficients) can only be applied if the left side is constant coefficient and the right side is a of a certain type.

If we want to solve an equation that doesn't have a constant coefficient left

$$\text{e.g. } x^2 y'' + xy' - 4y = e^x,$$

or doesn't have the correct sort of right hand side

$$\text{e.g. } y'' + y = \tan x,$$

we'll need another solution process.

Variation of Parameters

For the equation in standard form

$$\frac{d^2y}{dx^2} + P(x)\frac{dy}{dx} + Q(x)y = g(x),$$

suppose $\{y_1(x), y_2(x)\}$ is a fundamental solution set for the associated homogeneous equation. We seek a particular solution of the form

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x)$$

where u_1 and u_2 are functions we will determine (in terms of y_1 , y_2 and g).

$$y_c = c_1 y_1(x) + c_2 y_2(x)$$

This method is called **variation of parameters**.

Variation of Parameters: Derivation of y_p

$$y'' + P(x)y' + Q(x)y = g(x)$$

Set $y_p = u_1(x)y_1(x) + u_2(x)y_2(x)$

we have two unknowns u_1 and u_2 but only one equation, the ODE.
We'll introduce a 2nd equation.

Sub y_p into the ODE.

$$y_p = u_1 y_1 + u_2 y_2$$

$$y_p' = u_1 y_1' + u_2 y_2' + u_1' y_1 + u_2' y_2$$

Remember that $y_i'' + P(x)y_i' + Q(x)y_i = 0$, for $i = 1, 2$

Assume $u_1' y_1 + u_2' y_2 = 0$

This will be our 2nd equation.

So $y_p = u_1 y_1 + u_2 y_2$

$$y_p' = u_1 y_1' + u_2 y_2'$$

$$y_p'' = u_1' y_1' + u_2' y_2' + u_1 y_1'' + u_2 y_2''$$

$$y_p'' + P(x)y_p' + Q(x)y_p = g(x)$$

$$u_1' y_1' + u_2' y_2' + \underline{u_1 y_1''} + \underline{u_2 y_2''} + P(x)(\underline{u_1 y_1'} + \underline{u_2 y_2'}) + Q(x)(\underline{u_1 y_1} + \underline{u_2 y_2}) = g(x)$$

Collect u_1' , u_2' , u_1 and u_2

$$u_1' y_1' + u_2' y_2' + (y_1'' + P(x)y_1' + Q(x)y_1)u_1 \\ + (y_2'' + P(x)y_2' + Q(x)y_2)u_2 = g(x)$$

$$y_i'' + P(x)y_i' + Q(x)y_i = 0 \quad \text{for } i=1, \text{ or } 2$$

This reduces to

$$u_1' y_1' + u_2' y_2' = g(x)$$

We have the system of equations

$$u_1' y_1 + u_2' y_2 = 0$$

$$u_1' y_1' + u_2' y_2' = g(x)$$

We'll solve this using Cramer's rule.

$$\begin{bmatrix} y_1 & y_2 \\ y_1' & y_2' \end{bmatrix} \begin{bmatrix} u_1' \\ u_2' \end{bmatrix} = \begin{bmatrix} 0 \\ g \end{bmatrix}$$

Matrix
for the
Wronskian
of y_1, y_2

$$\text{Let } W_1 = \begin{vmatrix} 0 & y_2 \\ g & y_2' \end{vmatrix} = -g y_2$$

$$\text{and } W_2 = \begin{vmatrix} y_1 & 0 \\ y_1' & g \end{vmatrix} = y_1 g$$

$$\text{Let } W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \text{ the Wronskian of } y_1, y_2$$

$$u_1' = \frac{w_1}{w} = \frac{-g y_2}{w}$$

$$u_2' = \frac{w_2}{w} = \frac{y_1 g}{w}$$

$$u_1 = \int \frac{-g(x) y_2(x)}{w} dx \quad \text{and}$$

$$u_2 = \int \frac{g(x) y_1(x)}{w} dx$$

Example:

Solve the ODE $y'' + y = \tan x$.

$$y = y_c + y_p$$

$$\text{Find } y_c : y_c'' + y_c = 0$$

$$\text{Charact eqn } m^2 + 1 = 0$$

$$m^2 = -1 \Rightarrow m = \pm i$$

$$\alpha = 0, \beta = 1$$

$$y_1 = \cos x, \quad y_2 = \sin x$$

$$y_c = c_1 \cos x + c_2 \sin x$$

Using Variation of parameters, we need

W and g .

$$g(x) = \tan x, \quad W = \begin{vmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{vmatrix}$$

$$= \cos^2 x + \sin^2 x = 1$$

$$y_1 = \cos x, \quad y_2 = \sin x$$

$$u_1 = \int \frac{-gy_2}{W} dx = \int \frac{-\tan x \sin x}{1} dx$$

$$= - \int \frac{\sin^2 x}{\cos x} dx = - \int \frac{1 - \cos^2 x}{\cos x} dx$$

$$= \int (\cos x - \sec x) dx$$

$$u_1 = \sin x - \ln |\sec x + \tan x|$$

$$u_2 = \int \frac{g(x) y_1(x)}{W} dx = \int \frac{\tan x \cos x}{1} dx$$

$$u_2 = \int \sin x dx = -\cos x$$

$$y_1 = \cos x \quad y_2 = \sin x$$

$$y_p = u_1 y_1 + u_2 y_2$$

$$= (\sin x - \ln|\sec x + \tan x|) \cos x - \cos x \sin x$$

$$= \cancel{\sin x \cos x} - \cos x \ln|\sec x + \tan x| - \cancel{\cos x \sin x}$$

$$y_p = -\cos x \ln|\sec x + \tan x|$$

The general solution

$$y = C_1 \cos x + C_2 \sin x - \cos x \ln |\sec x + \tan x|$$

$$y'' + y = \tan x$$