

Lecture 18: Hamiltonian cycles


October 19, 2021

Kennesaw State University

1 Some examples

1.1 Knight's tours

A knight is a chess piece that (on the 8×8 chessboard) moves by taking two steps in one direction, turning 90° , and taking one more step. We are all graph theorists here, and so for us this means that a knight moves along the edges of the 64-vertex graph in the first diagram below. (For clarity, I have highlighted the 8 edges out of the typical vertex.)




The oldest Hamiltonian cycle problem in history is finding a **closed knight's tour** of the chessboard: the knight must make 64 moves to visit each square once and return to the start. That's exactly a Hamiltonian cycle in the graph we just drew. One solution is shown in the second diagram above.

We will not try to solve the 8×8 problem today. However, we will think about the problem for some smaller boards, and see what happens there!

1.2 The Icosian game

The problem of finding Hamiltonian cycles is named after William Rowan Hamilton, an Irish mathematician who was a big fan of the dodecahedron (the 3D shape whose sides are twelve pentagons). The graph below is the "skeleton graph" of the dodecahedron: the graph whose vertices are the corner points of the dodecahedron, and whose edges are the literal physical edges between those corner points.



¹This document comes from the Math 3322 course webpage: <http://facultyweb.kennesaw.edu/mlavrov/courses/3322-fall-2021.php>

Hamilton's "Icosian² game" was a puzzle to find a Hamiltonian cycle of this graph. The graph is small enough that we can try to do this by hand without fancy theorems. Usually the following approach will work:


1. Draw a few edges of the cycle as the fancy strikes you. (It's possible to make a mistake here, but unlikely.)
2. Every vertex has degree 2 in the cycle, but degree 3 in this graph. So if we've decided that edges uv and vw are part of the cycle, then the third edge out of v cannot be part of the cycle; get rid of it.
3. Similarly, if we've already eliminated one edge out of a vertex, it means that the cycle must use the other two edges through that vertex.
4. Finally, we should avoid edges that would result in the cycle looping back on itself too early (before it visits all 20 vertices).

If steps 2–4 don't give us enough information to solve for the rest of the cycle, then we can guess again.

As you can see, this is not a systematic approach: there's lots of guessing involved! The truth is that there's no systematic approach to finding Hamiltonian cycles that doesn't use brute force.

1.3 The Petersen graph

Does the Petersen graph (first diagram below) have a Hamiltonian cycle?



There's a way of thinking about this question which will be useful to us in future arguments. If the Petersen graph has a Hamiltonian cycle, then it should be possible to find a different drawing of the same graph in which all 10 vertices are arranged around the boundary of a circle, with adjacent edges connected (second diagram above).

The Petersen graph is 3-regular, so each vertex has a third edge out of it. But we don't know which edges to draw yet because we don't know if such a diagram exists: that depends on whether the Petersen graph has a Hamiltonian cycle or not.

The Petersen graph has an unusual property: it has no cycles of length 3 or 4. (Check this! It's enough to check that none of the neighbors of a vertex are connected, and that no two vertices share a neighbor.) So our hypothetical second diagram of the Petersen graph must also have this property.

²It might be confusing why it's called "icosian" when it's played on the dodecahedron; the icosahedron is a different shape with 20 triangular faces. Well, "icosa-" is the Greek prefix that means 20, and the graph we're looking at has 20 vertices.

In particular, from each vertex, there are only three possible candidates from the third edge: it must go either to the opposite vertex or to its neighbors. (Also shown in the second diagram.) Otherwise, we get a cycle of length 3 or 4.


It does not work to connect every vertex to its opposite. That gets us the third diagram, which definitely has cycles of length 4. So there's some vertex v that's *not* connected to its opposite vertex w . This is shown in the fourth diagram above, where v is the top vertex and w is the bottom vertex. But now, all three options for w 's third edge are bad: it cannot be v (that would make v have degree 4) and it cannot be either of v 's neighbors (that would create a cycle of length 4).

So actually, the diagram we want cannot exist, and the Petersen graph is not Hamiltonian!

2 Necessary conditions

The argument for the Petersen graph just now was long and complicated. Fortunately, in many cases, we can rule out a Hamiltonian cycle much more easily. Consider the following examples; some of these come from other situations we've looked at in class, and others are knight's graphs for smaller chessboards.

I recommend stopping to think of a reason for each of these graphs why it is not Hamiltonian.



Here's what we can say:

1. In the first graph (the 3×3 knight's graph) there is no way to get to the middle vertex, so there can't be a Hamiltonian cycle. A general principle: a graph that is not connected cannot be Hamiltonian.
2. In the second graph, some vertices have degree 1. There is no way a cycle can both enter and exit those vertices. A general principle: a Hamiltonian graph must have minimum degree at least 2.
3. In the third graph, there is a cut vertex. Hamiltonian graphs have no cut vertices, because for every two vertices v, w , there is a cycle containing both v and w (we can use a Hamiltonian cycle for every pair of vertices).
4. In the fourth graph, some vertices have degree 2, which is not a problem by itself. However, if a vertex has degree 2, both of its edges must be part of every Hamiltonian cycle. And if we select both edges out of every degree-2 vertex in this graph, we end up with two 4-cycles instead!
5. The fifth graph (the 5×5 knight's graph) is bipartite, as is any other knight's graph. In the usual black-and-white coloring of a chessboard, every move takes the knight to a square of a different color. In particular, along a Hamiltonian cycle, the colors should alternate black and white.

However, the 5×5 chessboard has 12 squares of one color and 13 of the other. These can't alternate black and white, because there's not an equal number of them.

A general principle: a bipartite graph in which the two parts are different sizes cannot be Hamiltonian.

Surprisingly, all five arguments are special cases of one result!

We say that a graph G is **tough** if, for any $k \geq 1$, deleting k vertices from G results in at most k connected components.

Lemma 2.1. *For all $n \geq 3$, the cycle graph C_n is tough.*

Proof. Deleting a single vertex from C_n leaves a path P_{n-1} , which is connected. Deleting a vertex from P_{n-1} leaves either a single path (if we delete from the endpoints) or two shorter paths (if we delete from the middle).

From then on, our graph will be a collection of paths. Deleting another vertex can, at most, break a path into two paths, increasing the number of components by one. Deleting k vertices total can leave at most k path components. \square

Corollary 2.2. *All Hamiltonian graphs are tough.*

Proof. Every n -vertex Hamiltonian graph G contains a copy of C_n as a subgraph. Deleting k vertices from G will leave that subgraph in at most k pieces, by the lemma above. The edges that G has that the Hamiltonian cycle does not can only reduce the number of components. \square

In each of the cases above, we can find a set of k vertices whose deletion leaves more than k components. (See if you can do this in all cases.) Therefore they are not tough—and therefore they are not Hamiltonian.

This is a necessary condition: to be Hamiltonian, a graph must be tough. However, it is not sufficient. It takes some casework, but we can check that the Petersen graph is tough, even though (as we saw earlier) it is not Hamiltonian.

3 Sufficient conditions

The prettiest condition that *does* guarantee Hamiltonian cycles in a graph comes from the Bondy–Chvátal theorem, which I like to call the “The magic was inside you all along” theorem.

Theorem 3.1 (Bondy–Chvátal). *In a graph G with n vertices, suppose s and t are two non-adjacent vertices with $\deg(s) + \deg(t) \geq n$.*

If $G + st$ (the graph we obtain from G by adding edge st) has a Hamiltonian cycle, then so does G .

Proof. Take a Hamiltonian cycle in $G + st$. If edge st is not part of that cycle, we're done: it also exists in G . If edge st is part of that cycle, then deleting that edge leaves an $s - t$ path in G that visits every single vertex: $(s = v_1, v_2, \dots, v_{n-1}, v_n = t)$

Moreover, because $\deg(s) + \deg(t) \geq n$, there must be many other edges from s or t in the graph. Let's take a closer look at those edges.



Let S be the set of edges $v_i v_{i+1}$ on the path in which v_{i+1} (the endpoint closer to t) is adjacent to s . These are highlighted in red in the first diagram above (for an example set of edges out of s). Because we get one edge in S for every neighbor of s , we know that $|S| = \deg(s)$.

Similarly, let T be the edges $v_i v_{i+1}$ on the path in which v_{i+1} (the endpoint closer to s) is adjacent to t . These are highlighted in blue in the second diagram; as before, $|T| = \deg(t)$.

Because $|S| + |T| = \deg(s) + \deg(t) \geq n$, but the path only has $n - 1$ edges, there must be overlap! Some edge $v_i v_{i+1}$ must be both in S and in T .



In this case, $(s = v_1, v_2, \dots, v_i, t = v_n, v_{n-1}, \dots, v_{i+1}, s)$ is a Hamiltonian cycle. □

Essentially, the Bondy–Chvátal theorem says that edges like st don't help create a Hamiltonian cycle if one wasn't already there.

To apply this theorem, we use the following logic. Take any graph G , then add every single edge that the theorem applies to. (The resulting graph is called the **Bondy–Chvátal closure** of G .) If the result is Hamiltonian, then the original graph is also Hamiltonian. However, it might be easier to find a Hamiltonian cycle in the new graph, because it has more edges to work with.

Corollary 3.2 (Dirac's theorem). *If G has $n \geq 3$ vertices and minimum degree $\delta(G) \geq \frac{1}{2}n$, then G is Hamiltonian.*

Proof. Here, if we take the closure of G , we end up with K_n : for any two non-adjacent vertices s and t , $\deg(s) + \deg(t) \geq \frac{1}{2}n + \frac{1}{2}n = n$. Since K_n is definitely Hamiltonian, G must also be Hamiltonian. □